


**Definition of the pnictogen bond**

Journal:	<i>Pure and Applied Chemistry</i>
Manuscript ID	PAC-REC-2020-1002.R4
Manuscript Type:	Recommendation
Date Submitted by the Author:	17-Oct-2022
Complete List of Authors:	<p>Resnati, Giuseppe; Politecnico di Milano Dipartimento di Chimica Materiali e Ingegneria Chimica Giulio Natta, Chemistry, Materials, Chemical Engineering "Giulio Natta"</p> <p>Bryce, David</p> <p>Desiraju, Gautam; Indian Institute of Science, Solid State &amp; Structural Chemistry Unit</p> <p>Frontera, Antonio; Universitat de les Illes Balears Departament de Química</p> <p>Krossing, Ingo; University of Freiburg</p> <p>Legon, Anthony; university of bristol, School of Chemistry,</p> <p>Metrangolo, Pierangelo; Politecnico di Milano, Dipartimento di Chimica, Materiali e Ingegneria Chimica "Giulio Natta"</p> <p>Nicotra, Francesco; University of Milano-Bicocca, Department of Bioscience and Biotechnology</p> <p>Rissanen, Kari; University of Jyväskylä</p> <p>Scheiner, Steve; Utah State University, Department of Chemistry and Biochemistry</p> <p>Terraneo, Giancarlo; Politecnico di Milano, Dipartimento di Chimica, Materiali e Ingegneria Chimica "Giulio Natta"</p>
Keywords:	pnictogen bond, IUPAC Organic and Biomolecular Chemistry Division, IUPAC Physical and Biophysical Chemistry Division, nomenclature, noncovalent interactions, self-assembly
Author-Supplied Keywords:	

SCHOLARONE™  
Manuscripts

1

Giuseppe Resnati\*, David L. Bryce, Gautam R. Desiraju, Antonio Frontera, Ingo Krossing, Anthony C. Legon, Pierangelo Metrangolo, Francesco Nicotra, Kari Rissanen, Steve Scheiner, and Giancarlo Terraneo

# Definition of the pnictogen bond

## (IUPAC Recommendations 202?)

<http://doi.org/10.26434/chemrxiv-2016-001-2-300>  
Received

**Abstract:** This recommendation proposes a definition for the term “pnictogen bond”: the term pnictogen bond designates a subset of the attractive interactions between an *electrophilic* region on a *pnictogen* atom in a *molecular entity* and a *nucleophilic* region in another, or the same, molecular entity.

**Keywords:** pnictogen bond, IUPAC Organic and Biomolecular Chemistry Division, IUPAC Physical and Biophysical Chemistry Division, nomenclature, noncovalent interactions, self-assembly, supramolecular chemistry.

### CONTENTS

#### 1. PREFACE

#### 2. DEFINITION

2.1 List of Some Common Pnictogen Bond Donors and Acceptors

2.2 List of Features

#### 3. ISSUES ON PNICTOGEN BOND

#### 4. TERMS

#### MEMBERSHIP OF SPONSORING BODIES

#### REFERENCES

**Article note:** This work was started under project 2016-001-2-300: Categorizing Chalcogen, Pnictogen, and Tetrel Bonds, and Other Interactions Involving Groups 14-16 Elements.

**\*Corresponding author: Giuseppe Resnati**, NFMLab, Dept. Chemistry, Materials, and Chemical Engineering “Giulio Natta”, Politecnico di Milano, Via Mancinelli 7, I-20131 Milano, Italy, e-mail: giuseppe.resnati@polimi.it. <http://orcid.org/0000-0002-0797-9296>

**David L. Bryce:** Dept. Chemistry and Biomolecular Sciences & Centre for Catalysis Research and Innovation, University of Ottawa, 10 Marie Curie Private, Ottawa, Ontario K1N 6N5, Canada

**Gautam R. Desiraju:** Solid State and Structural Chemistry Unit, Indian Institute of Science, Bangalore, 560 012, India

**Antonio Frontera:** Dept. Química, Universitat de les Illes Balears, Crta de Valldemossa km 7.5, 07122 Palma de Mallorca (Balears), Spain

**Ingo Krossing:** Inst. Inorganic and Analytical Chemistry, Albert-Ludwigs-Universität Freiburg, Albertstrasse 21, D-79104 Freiburg, Germany

**Anthony C. Legon:** School of Chemistry, University of Bristol, Bristol BS8 1TS, Avon, UK

**Francesco Nicotra:** Dept. Biotechnologies and Biosciences, University of Milano Bicocca, Piazza della Scienza, 2, I-20126 Milano, Italy

**Kari Rissanen:** Dept. Chemistry, University of Jyväskylä, POB 34, Jyväskylä 40014, Finland

**Steve Scheiner:** Dept. Chemistry and Biochemistry, Utah State University, Logan, UT 84322-0300, USA

**Giancarlo Terraneo and Pierangelo Metrangolo:** SBNLab, Dept. Chemistry, Materials, and Chemical Engineering “Giulio Natta”, Politecnico di Milano, Via Mancinelli 7, I-20131 Milano, Italy

## 1. Preface

This recommendation proposes a definition for the term “pnictogen bond” (PnB) and suggests it to be used to designate a subset of the inter- and intramolecular interactions formed by pnictogen atoms (i.e., atoms of Group 15 elements) bonded in a molecular entity [1]. The alternative spelling ‘pnicogen bond’ can also be used to designate these bonds, the latest IUPAC Recommendations on Inorganic Chemistry stating that pnictogen is IUPAC-approved and the alternative spelling pnicogen is also used [2].

In this recommendation the use of the terms donor, acceptor, and of the derived expressions is consistent with the use of the same terms for the hydrogen bond (HB) [3, 4], the halogen bond (HaB) [5], and the chalcogen bond (ChB) [6], three interactions that present similarities with the PnB. The respective IUPAC recommendations specify that the HB donor, the HaB donor, and the ChB donor are the molecular entities containing the electrophilic [1] hydrogen, halogen, and chalcogen atoms, respectively. Hereinafter, the PnB donor is the molecular entity containing the electrophilic pnictogen atom, the PnB donor-atom is the electrophilic pnictogen atom, and the PnB donor-region is the electrophilic region on the PnB donor-atom. The PnB acceptor is the molecular entity functioning as the nucleophile [1], similar to the terminology used for the HB, HaB, and ChB. In other words, for HB, HaB, ChB, and PnB, the interaction donor and the interaction acceptor are the acceptor of electron density and the donor of electron density, respectively. The terms PnB acceptor-atom and PnB acceptor-region (as well as the terms HB donor-atom/region, HaB donor-atom/region, and ChB donor-atom/region) are used in this recommendation according to consistent meanings.


Covalently bonded atoms have an anisotropic distribution of the electron density and present regions of higher and lower electron density at their outer zones. At these regions, the molecular surface electrostatic potential is frequently negative and positive, respectively [7-9], and attractive interactions with regions of opposite electrostatic potential can be formed.

Hydrogen atoms bonded to atoms, or groups, more electronegative than themselves are partially positive and this is at the origin of the HB [3, 4]. The surface electrostatic potential at hydrogen outer regions is non-uniform [10] and this is related to HB directionality at the hydrogen atoms (the HB donor-atoms).

Halogen atoms forming one covalent bond can function as electron-rich atoms (i.e., nucleophiles, Lewis bases [1]). This is the case, for instance, when halogen atoms donate electron density to, and attractively interact with, hydrogen atoms or metal cations [11-14] approaching the halogen preferentially at its lone pairs. Halogen atoms can also function as electron-poor atoms (i.e. electrophiles, Lewis acids [1]). This is the case, for instance, when they accept electron density from, and attractively interacting with, lone-pair possessing atoms and anions [15] approaching the halogen approximately on the extension of, and opposite to, the covalent bond formed by the halogen (Scheme 1-left). The halogen lone pairs and the region opposite to the covalent bond at halogen ( $\sigma$ -hole) are the positions of preferred formation of short contacts with Lewis acids and bases, respectively, as they correspond to the regions of higher and lower electron density where the electrostatic potential is typically negative and positive, respectively. The IUPAC recommendation defining the halogen bond [5] acknowledges that an electrophilic behavior can be shown by halogen atoms in quite different moieties [1], including molecular entities wherein halogen atoms form more than one covalent bond [16, 17]. This recommendation establishes that a halogen bond occurs when there is evidence of a net attractive interaction between an electrophilic region associated with a halogen atom in a molecular entity and a nucleophilic region in another, or the same, molecular entity.

Similarly, chalcogen atoms can behave as both nucleophiles and electrophiles (Lewis bases and acids, respectively). The nucleophilic behaviour of chalcogen atoms is most common for oxygen, the most electronegative [1, 18] and least polarizable [1, 19] chalcogen, e.g., oxygen can frequently function as a hydrogen bond acceptor-atom. Other elements of Group 16 can behave as nucleophiles, e.g., when donating electron density to a halogen or hydrogen atom [20, 21]. The electrophilic behaviour of chalcogen atoms is common for the less electronegative and more polarizable elements of the group, these elements frequently form attractive interactions with neutral or negatively charged nucleophiles (typically amine or pyridine nitrogen atoms or halide anions) [22]. The two classes of interactions described

3



20  
21  
22  
23

**Scheme 1.** Drawing evidencing the preferred location of electrophilic regions in monovalent halogens (left), divalent chalcogens (mid), and trivalent pnictogens (right). These electrophilic regions ( $\sigma$ -holes) are on the extension of the covalent bonds formed by the respective atoms.

24  
25  
26  
27  
28  
29  
30  
31  
32  
33


above differ in their electronic features and their directionalities relative to the covalent bonds formed by the chalcogen atom. Chalcogen atoms forming two single covalent bonds are approached by electrophiles after a trajectory roughly  $20^\circ$  from the perpendicular to the plane through the two covalent bonds as in these positions the presence of the lone-pairs tends to cause the existence of excess electron density and negative electrostatic potential. Differently, the approach of nucleophiles to the chalcogen atoms is preferentially on the extensions of the two covalent bonds, the regions named  $\sigma$ -holes where the electron density is depleted and the electrostatic potential is frequently positive (Scheme 1-mid). An IUPAC recommendation [6] recently recognized the general ability of chalcogen atoms, hypervalent [23] chalcogen atoms included [24], to interact with nucleophiles and proposed to name as chalcogen bonds (ChBs) only those attractive interactions wherein the chalcogen atom is the electrophile.

34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48

Elements of Group 15 can behave as electron-rich (nucleophilic) atoms and form attractive interactions with electron-poor (electrophilic) partners. This is the case, for instance, when pnictogen atoms act as HB [3, 4] or HaB [5, 15] acceptors (Scheme 2-top). This nucleophilic behaviour is most common for nitrogen, the element with the highest electronegativity and the lowest polarizability in the Group, in analogy with what is described above for chalcogen atoms. It has long been known that Group 15 atoms can additionally function as electron-poor (electrophilic) atoms, i.e., they can attractively interact with electron-rich (nucleophilic) atoms (Scheme 1-right, Scheme 2-bottom). Antimony, thanks to its low electronegativity and high polarizability, can function as a particularly strong Lewis acid. For example, antimony pentachloride is the standard Lewis acid in the time honoured Gutmann scale [32] of Lewis basicity and antimony pentafluoride is a reference compound to identify Lewis superacids [33]. Interactions formed by Group 15 elements in the two classes of interactions described above differ in their electronic features and differ also in their directionality relative to the covalent bonds formed by the pnictogen atom as the regions of preferential approach of electrophiles or nucleophiles are those with negative or positive surface electrostatic potential, respectively.

49  
50  
51  
52  
53  
54  
55  
56  
57

Acknowledging an established usage in the chemical literature [34], this recommendation proposes that the term “pnictogen bond” (PnB) is used to designate uniquely the specific subset of the inter- and intramolecular interactions wherein Group 15 elements act as electrophile [35] and wherein separations, energies, and/or other experimental/theoretical parameters significantly differ from those of corresponding covalent/ionic bonds. A short definition of the term is given at the beginning of the next section, followed by explanatory examples in the form of a not exhaustive list of some common PnB donors and acceptors. A list of experimental and/or theoretical features follows; although not comprehensive, these



**Scheme 2.** Typical cases where: nitrogen functions as nucleophile and forms HBs [25] or HaBs [26] on interacting with an atom (El) with electrophilic regions (top); bottom, nitrogen [27], phosphorus [28], or antimony [29, 30] function as atoms with electrophilic regions and form PnBs on interaction with nucleophiles (Nu) (bottom). Arrows indicate the approach direction of El/Nu. In the two left cases of electrophilic pnictogens, the regions with positive electrostatic potential and electrophilic character (typically named  $\pi$ -holes [7, 28, 31]) are approximately above and below the plane of the reported moiety [1]; Nu approaches N and P atoms at right angle to the plane of the moiety. In the two right cases of electrophilic pnictogens, the regions with positive electrostatic potential and electrophilic character (typically named  $\sigma$ -holes [7-9, 28, 31]) are approximately on the extension of, and opposite to, a covalent bond at Sb; Nu approaches this atoms after directions approximately coaxial with this bond.

features can be used as evidence for the presence of a pnictogen bond. This recommendation is concluded by a discussion of some important issues on the PnB, in order to provide the reader with qualitative information for a correct use of the term. This recommendation runs purposefully side by side with the recommendations for the terms halogen bond and chalcogen bond [5, 6]. Similarities in the overall logical flow and in some sentences of these three recommendations are pursued in order to develop a consistent set affording a systematic terminology for the different interactions which are formed by different elements but possess similarities in geometric and electronic features.

In order to better outline and contextualize the pnictogen bond defined in this recommendation, it might be useful to mention and briefly discuss some other terms which are used in the literature. The term secondary bond [36] is used to indicate a set of different interactions formed by p-block elements and sharing the common feature that the interaction is collinear with a covalent bond at the p-block element. The terms  $\sigma$ -hole bond and  $\pi$ -hole bond [7-9, 28] are employed as umbrella terms for interaction sets wherein the shared feature is that the nucleophile approaches the electrophile collinear with a covalent bond formed by the electrophile or orthogonal to a molecular portion of the electrophile, respectively.

## 2. Definition

### **pnictogen bond (PnB)**

weak attractive interaction between an *electrophilic* region on a *pnictogen* atom in a *molecular entity* (wherein the *pnictogen* is involved in other stronger bonds) and a *nucleophilic* region in another, or the same, *molecular entity*.

Note 1: A typical pnictogen bond is denoted by the three dots in R–Pn⋯A, wherein R is collectively representing the residues covalently bonded to the pnictogen atom Pn, Pn (the PnB donor-atom) is any atom (possibly hypervalent [37]) of Group 15 having at least one electrophilic (electron-poor) region (the PnB donor-region) involved in a bond with A, R–Pn (the PnB donor) is a molecular entity containing the electrophilic pnictogen atom, and A (the PnB acceptor) is a molecular entity having at least one nucleophilic (electron-rich) region (the PnB acceptor-region).

Note 2: An electrophilic region on a pnictogen atom in a molecular entity and a nucleophilic region in another or the same molecular entity can form bonds spanning a range of separations, energies, and other experimental or theoretical parameters; the use of the term pnictogen bond has to be limited to cases where separations, energies, and/or other experimental/theoretical parameters significantly differ from those of corresponding covalent/ionic bonds (see onwards).

Note 3: The polar character of the pnictogen atom in the interaction (i.e., the electrophilic role in pnictogen bonded systems or its nucleophilic role in the other systems) is established by considering the physical and chemical properties of the starting molecular entities before the interaction formation and/or of the system assembled by the interaction (see onwards).

Note 4: Bonds wherein a pnictogen atom acts as the nucleophile and those wherein its electrophilic role cannot be identified must not be named pnictogen bonds.

Note 5: Pnictogen atoms can concurrently form one or more than one pnictogen bond [38]; the PnB donor and/or PnB acceptor can be charged molecular entities thus forming charge assisted PnBs.

### **2.1. On Common Pnictogen Bond Donors and Acceptors**

The elements of Group 15 form bonds with many other elements so that pnictogens can produce a very wide diversity of functional groups characterized by quite different properties. A list of common PnB donors containing functional groups [1] whose pnictogen atoms are particularly prone to form PnBs would thus be either very long or highly incomplete. In order to provide a quick but fairly comprehensive identification of such functional groups, we report below some selected atoms and groups which effectively promote the PnB donor ability of a pnictogen atom when covalently bonded to it.

A non-comprehensive list of atoms and groups promoting the PnB formation includes:

- fluorine, chlorine, and bromine atoms (as in homocrystals of dichloro-1,1-dimethylethylstibane [39], dichloro(phenyl)stibane [40], and dibromo(4-methylphenyl)stibane [41] wherein halogen atoms are the nucleophile);
- oxygen groups, namely alkoxy, aryloxy, acyloxy, or oxo groups (as in nitroalkyl and nitroaryl derivatives [27, 42], di(quinoline-7-yl)cyclohexylphosphonite [43], (-)-(1*S*,5*S*,7*S*)-2,4,10-trioxa-1,3,5,7-tetraarsadamantane ((-)-arsenicin A, [44]), 2,4,6,8-tetramethyl-1,3,5,7,2,4,6,8-tetraoxatetraarsocane [45], tris(2,2-dimethylpropanoato)arsenic) [46] wherein oxygen atoms are the nucleophile);
- cyano and perfluoroalkyl groups (as in dimethylarsanecarbonitrile [47], tris(pentafluoroethyl)bismuthane [48] wherein nitrogen and fluorine atoms are the nucleophile, respectively);
- positively charged heteroaromatic group (as in (diphenylbismuthanyl)trimethylphosphonium chloride [49], 2,2'-(chlorophosphanediy)bis(1,3-diisopropyl-4,5-dimethyl-1*H*-imidazol-3-ium)

trifluoromethanesulfonate [50], 2-(dichlorostibanyl)-1,3-diisopropyl-4,5-dimethyl-1*H*-imidazol-3-ium trifluoromethanesulfonate [51], 2-(dibromostibanyl)-1,3-bis(2,6-diisopropylphenyl)-1*H*-imidazol-3-ium tetrabromoaluminate [52] wherein the respective anions are the nucleophile).

A non-comprehensive list of PnB acceptors A includes:

- lone-pair possessing atoms (e.g., N atom of pyridines or amines, O atom of ethers or carbonyl groups);
- $\pi$  systems (e.g., double or triple bonds between carbon atoms, arene moieties);
- anions (e.g., when halide anions or polyatomic oxyanions interact with the neutral phosphorus of 1-(4-carboxymethyl-benzyl)-1,3,5-triaza-7-phosphaadamantan-1-ium bromide [53] or the neutral antimony of 5-ethyl-2,3,7,8,14,15-hexamethyl-5*H*-5,10-[1,2]benzenodibenzo[b,e][1,4]phosphastibinin-5-ium iodide and its bromide analogue [54]).

## 2.2. List of Features of Pnictogen Bond

The evidence for the occurrence of a PnB may be experimental or theoretical, or better, a combination of both. A non-exhaustive list of features that are useful as indications for the PnB presence is reported below. The greater is the number of satisfied features, the more reliable the characterization of an interaction as a PnB.

In a typical pnictogen-bonded adduct R–Pn $\cdots$ A:

- in neutral adducts, the interatomic distance between the PnB donor-atom Pn and the PnB acceptor-atom A is marginally smaller than the sum of the respective van der Waals radii [55] and substantially longer than the sum of covalent radii [56];
- the PnB acceptor A tends to approach the donor-atom Pn either along the extension of one of the  $\sigma$  covalent bonds at Pn and opposite to the bond (forming the so-named  $\sigma$ -holes bonds [7-9, 28, 31]) or perpendicular to the plane formed by three coplanar covalent bonds at Pn (forming the so-named  $\pi$ -holes bonds [7, 28, 31]); if the former geometry is observed, deviation from the linear extension is frequently more pronounced in PnBs than in ChBs and HaBs [8];
- the pnictogen atom interacts with a lone-pair orbital approximately along the orbital axis and with a  $\pi$  region perpendicular to the  $\pi$  bond plane [28];
- the R–Pn covalent bond opposite to the PnB tends to be longer in the pnictogen bonded adduct than in the unbonded R–Pn moiety [57];
- with a given acceptor A, the PnB strength typically decreases when the electronegativity of Pn increases and the electron withdrawing ability of R in R–Pn decreases [31, 34, 35, 58-60];
- the analysis of the electron density topology usually shows a bond path and a bond critical point [23] between Pn and A;
- the infrared absorption and Raman scattering observables of both R–Pn and A are affected by PnB formation [57, 61-63];
- the UV–vis absorption bands of the PnB donor chromophore change upon PnB formation [64];
- the Pn $\cdots$ A PnB formation typically affects the nuclear magnetic resonance observables (e.g., chemical shift values, quadrupolar and spin-spin coupling constants) of nuclei in both R–Pn and A [57, 65-67].

## 3. Issues on Pnictogen Bond

Group 15 elements range across elements with quite different characteristics spanning from non-metals to metals, and the bonds landscape of derivatives of the lightest pnictogens can differ substantially from that of the heaviest ones. Nitrogen, the most electronegative pnictogen [18], is the least prone to function as electrophile. This element typically affords the weakest and longest PnBs [7, 8], the separation of involved atoms being usually close to or slightly smaller than the sum of van der Waals radii of involved atoms. Electrophilicity and PnB donor ability increase moving from phosphorus to arsenic and antimony, namely when electronegativity decreases [35, 64, 66-68]. Bismuth, the least electronegative pnictogen, is the most disposed to act as electrophile and to expectedly form strong and short PnBs. Indeed, when bismuth derivatives interact with donors of electron density, the interaction

energy and separation are frequently close to or slightly longer than those of the corresponding covalent bonds. These contacts are typically understood as coordinative interactions [68] and have not to be named pnictogen bonds, this latter term applying only to longer and weaker contacts. Also some contacts between nucleophiles and trivalent or pentavalent derivatives of antimony, or less frequently of arsenic, are characterized by energies and separations similar to those of corresponding covalent bonds. This is the case, for instance, for some bonds formed by Cl or Br with BiCl<sub>3</sub>, SbCl<sub>5</sub>, AsCl<sub>5</sub>, or their fluorinated analogues, which can be as short as Bi/Sb/As–Cl/F covalent bonds in the starting molecular entities; when so, these bonds must not be named pnictogen bonds.

The bonds geometry at the pnictogen atom in the isolated PnB donor may undergo a change on PnB formations and deformation tends to increase with the PnB strength. In an adduct wherein a group 15 atom functions as the electrophile, the short contact formed with the nucleophile is a pnictogen bond limited to cases where the geometry of the PnB donor undergoes a minor rearrangement on bond formation.

In the Preface of this recommendation the PnB was introduced by adopting a viewpoint based on electrostatics, but calculations indicate that, similar to HaB and ChB, additional effects also contribute to the PnB formation. The seminal statement on the HaB "The *attractive nature* of the interaction is mainly due to electrostatic effects, but polarization, charge-transfer, and dispersion contributions all play an important role in causing interpenetration of van der Waals volumes" [69], holds also for the PnB [70-73]. The relative relevance of these and other components (the covalent one included) depends, among others, on the nature of the involved atoms and of the residues covalently bonded to them.

A list is reported below of exemplary cases where the pnictogen atom forms an attractive interaction, via an electrophilic region, with a nucleophilic region in another, or the same, molecular entity and the use of the term pnictogen bond for the resulting bond is consistent with the definition given above (Scheme 3). Another list follows of exemplary cases wherein the pnictogen atom functions as the nucleophile (Lewis base) and the use of the term pnictogen bond is inappropriate (Scheme 4). The naming of interactions sketched in Schemes 3 and 4 is consistent with the relevant IUPAC definitions [3-6], with the established ability of pnictogen atoms to act as both nucleophiles [15, 74-76] and electrophiles [32-35], and with the different terms traditionally employed to indicate the variety of bonds formed by pnictogen atoms [36]. Importantly, the use of the term pnictogen bond for interactions in Scheme 3 is consistent with the whole history of the term, from the very initial employments [77,78] to the most recent systematizations [79], going through its position in a taxonomy of chemical interactions [80].

Cases wherein the group 15 element forms a pnictogen bond (Scheme 3):

- the P···O interaction in the PCl<sub>3</sub>/H<sub>2</sub>O, PCl<sub>3</sub>/CH<sub>3</sub>OH, POCl<sub>3</sub>/H<sub>2</sub>O dimers [61-63] and the P···N interaction in di(quinolin-7-yl)cyclohexylphosphonite [43];
- the As···N interaction in dimethylarsanecarbonitrile [47] and the As···O interaction in (-)-2,4,10-trioxa-1,3,5,7-tetraarsadamantane [44];
- the Sb···Cl, Sb···Br, and Sb···S interactions in dichloro(phenyl)stibane [40] and its dibromo analogue [41] and in the adduct formed by SbCl<sub>3</sub> with 1,3,5-trithiane [82], respectively;
- the Bi···Se, Bi···Br, and Bi···Cl interactions in the cocrystal between BiCl<sub>3</sub> and triphenyl-λ<sup>5</sup>-phosphaneselenone [83], in crystalline di(2,4,6-trimethylphenyl)bromobismuthane [84] and (diphenylbismuthanyl)trimethylphosphonium chloride [49].

Cases wherein the interaction formed by the group 15 element is not a pnictogen bond (Scheme 4):

- the contacts observed in the solid between the NH group of 5,5-diethylpyrimidine-2,4,6-(1*H*,3*H*,5*H*)-trione and a pyrimidine nitrogen [74] or in the gas between HX (X=F, Cl, Br, I) and the phosphorus of phosphane [4, 75]; these interactions are hydrogen bonds (HBs);
- the attractive intermolecular interaction between the pnictogen atom in triphenylphosphane, triphenylarsane, and triphenylstibane and an iodine atom of 1,3,5-trifluoro-2,4,6-triiodobenzene in the corresponding co-crystals; these interactions are halogen bonds (HaBs) [76];
- the phosphorus-chalcogen contacts present in crystals of *N,N'*-{[diselanediy]bis(2,1-phenylene)}bis(methylene)}bis(*N*-propylpropan-1-amine) [85] and diisopropyl[6-(phenyltellanyl)-1,2-dihydroacenaphthylen-5-yl]phosphane [86]; these interactions are chalcogen bonds (ChBs);
- the contacts between nitrogen and tin in crystalline tris[2-(dimethylaminomethyl)phenyl]stannane derivatives [87] and the analogous short contacts between




8

1  
2  
3 nitrogen and germanium, or silicon, in crystalline tris[2-(aminomethyl)phenyl]germane [88], or  
4 silane [89], derivatives; in these interactions the electrophile is the 14 group element.  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60


For Peer Review Only

9




**Scheme 3.** Structural formulas of systems wherein the group 15 element forms PnBs (dotted line). Pnictogen bond separations (pm) and the normalized contacts (Nc) are reported close to the interactions. Normalized contacts for an interaction between atoms  $i$  and  $j$  is the ratio  $D_{ij}/(r_{vdW_i} + r_{vdW_j})$  where  $D_{ij}$  is the experimental or computed distance between  $i$  and  $j$  and  $r_{vdW_i}$  and  $r_{vdW_j}$  are the van der Waals radii [55] of  $i$  and  $j$ ; if the electron donor  $j$  is an anionic atom,  $r_{vdW_j}$  is substituted by  $rP_j$ , the Pauling ionic radius of anion atom  $j$  [81]. Normalized contacts are useful indicators because they allow for a more informative comparison of distances between different interacting atoms than absolute values of separations.


**Electrophile: Hydrogen atom  
(Hydrogen Bond, HB)**





**Electrophile: Group 17 Element  
(Halogen Bond, HaB)**



**Electrophile: Group 16 Element  
(Chalcogen Bond, ChB)**



**Electrophile: Group 14 Element**



**Scheme 4.** Structural formulas of systems wherein the bond between interacting moieties (dashed line) must not be named pnictogen bond.

## 4. Terms

Topical chemical terms used in this recommendation but missing to now an IUPAC definition are listed below. For these terms, citation are given to paper where the meaning of the term is the same as in this recommendation:

- $\pi$ -hole bond [7, 28, 64];
- $\sigma$ -hole bonds [7-9, 28, 64].

## Membership of sponsoring bodies

Membership of the Organic and Biomolecular Chemistry Division at the time of submission of this recommendation was as follows: President: N. E. Nifantiev (Russia); Past President: Francesco Nicotra (Italy); Secretary: S. Jarosz (Poland); Vice President: A. P. Rauter (Portugal); Titular Members: J. B. Harper (Australia), Y. Krishnan (USA), L. L. Mammino (South Africa), A. Marx (Germany), E. Uggerud (Norway), Z. Xi (China); Associate Members: P. Andersson (Sweden), F. Bai (China), G. Pandey (India), P. Scrimin (Italy), I. Shin (Korea), M. Yamashita (Japan); National Representatives: M. Donnard (France), M. Hegazy (Egypt), J. Honek (Canada), P. Kongsaree (Thailand), A. M. P. Koskinen (Finland), D. Milic (Serbia), E. D. Naydenova (Bulgaria), S. O. Okeniyi (Nigeria), R. Orru (Netherlands), S.-S. Sun (Taiwan). Membership of the Physical and Biophysical Chemistry Division at the time of submission of this recommendation was as follows: President: T. J. Wallington (United States); Past President: R. D. Weir (Canada); Secretary: A. Császár (Hungary); Vice President: P. Metrangolo (Italy); Titular Members: J. G. Frey (United Kingdom), F. Separovic (Australia), Z. Shuai (China), H. Tokoro (Japan), I. Voets (Netherlands), A. K. Wilson (USA); Associate Members: M. Fall (Senegal), J. L. Faria (Portugal), S.-J. Jeon (Korea), T. C. Kurtén (Finland), L. A. Montero-Cabrera (Cuba), V. Tsakova (Bulgaria); National Representatives: G. Ciric-Marjanovic (Serbia), M. Deyab (Egypt), L. C. Ngozi-Olehi (Nigeria), R. Orinakova (Slovakia), I. Vorotyntsev (Russia).

## References

- [1] P. Muller. *Pure Appl. Chem.* **66**, 1077–1184 (1994).
- [2] N. G. Connelly, T. Damhus, R. M. Hartshorn, A. T. Hutton. *Inorganic Chemistry IUPAC Recommendations 2005*, Royal Society of Chemistry, Cambridge, 2005, page 51. ISBN 0 85404 438 8.
- [3] E. Arunan, G. R. Desiraju, R. A. Klein, J. Sadlej, S. Scheiner, I. Alkorta, D. C. Clary, R. H. Crabtree, J. J. Dannenberg, P. Hobza, H. G. Kjaergaard, A. C. Legon, B. Mennucci, D. J. Nesbitt. *Pure Appl. Chem.* **83**, 1637–1641 (2011).
- [4] E. Arunan, G. R. Desiraju, R. A. Klein, J. Sadlej, S. Scheiner, I. Alkorta, D. C. Clary, R. H. Crabtree, J. J. Dannenberg, P. Hobza, H. G. Kjaergaard, A. C. Legon, B. Mennucci, and D. J. Nesbitt. *Pure Appl. Chem.* **83**, 1619–1636 (2011).
- [5] G. R. Desiraju, P. S. Ho, L. Kloo, A. C. Legon, R. Marquardt, P. Metrangolo, P. Politzer, G. Resnati, K. Rissanen. *Pure Appl. Chem.* **85**, 1711–1713 (2013).
- [6] C. B. Aakeroy, D. L. Bryce, G. R. Desiraju, A. Frontera, A. C. Legon, F. Nicotra, K. Rissanen, S. Scheiner, G. Terraneo, P. Metrangolo, G. Resnati. *Pure Appl. Chem.* **91**, 1889–1892 (2019).
- [7] P. Politzer, J. S. Murray. *ChemPhysChem* **21**, 579–588 (2020).
- [8] P. Politzer, J. S. Murray, T. Clark, G. Resnati. *Phys. Chem. Chem. Phys.* **19**, 32166–32178 (2017).
- [9] P. Politzer, J. S. Murray. *Crystals* **7**, 212 (2017).
- [10] J. S. Murray, P. Politzer. *J. Indian Inst. Sci.* **100**, 21–30 (2020).
- [11] G. Aullón, D. Bellamy, L. Brammer, E. A. Bruton, A. G. Orpen. *Chem. Commun.* 653–654 (1998).
- [12] C. B. Aakeröy, T. A. Evans, K. R. Seddon, I. Pálkó. *New J. Chem.* 145–152 (1999).
- [13] K. Molcanov, B. Kojic-Prodic, A. Meden. *Croat. Chem. Acta.* **82**, 387–396 (2009). CCDC no.: 695075.
- [14] J. Stierstorfer, K. R. Tarantik, T. M. Klapotke. *Chem. Eur. J.* **15**, 5775–5792 (2009). CCDC no.: 704307.
- [15] G. Cavallo, P. Metrangolo, R. Milani, T. Pilati, A. Priimagi, G. Resnati, G. Terraneo. *Chem. Rev.* **116**, 2478–2601 (2016).
- [16] G. Cavallo, J. S. Murray, P. Politzer, T. Pilati, M. Ursini, G. Resnati. *IUCrJ.* **4**, 411–419 (2017).
- [17] F. Heinen, E. Engelage, C. J. Cramer, S. M. Huber. *J. Am. Chem. Soc.* **142**, 8633–8640 (2020).
- [18] M. Rahm, T. Zeng, R. Hoffmann. *J. Am. Chem. Soc.* **141**, 342–351 (2019).
- [19] P. Schwerdtfeger, J. K. Nagle. *Molecular Phys.* **117**, 1200–1225 (2019).
- [20] H. D. Arman, R. L. Gieseking, T. W. Hanks, W. T. Pennington. *Chem. Commun.* **46**, 1854–1856 (2010).
- [21] A. Chand, D. K. Sahoo, A. Rana, S. Jena, H. S. Biswal. *Acc. Chem. Res.* **53**, 1580–1592 (2020).
- [22] P. Scilabra, G. Terraneo, G. Resnati. *Acc. Chem. Res.* **52**, 1313–1324 (2019).

- [23] V. I. Minkin. *Pure Appl. Chem.* **71** 1919–1981 (1999).
- [24] A. Franconetti, D. Quiñonero, A. Frontera, G. Resnati. *Phys. Chem. Chem. Phys.* **21**, 11313–11319 (2019).
- [25] T. W. Bell, A. B. Khasanov, M. G. B. Drew. *J. Am. Chem. Soc.* **124** 14092–14103 (2002).
- [26] P. Cardillo, E. Corradi, A. Lunghi, S. Valdo Meille, M. T. Messina, P. Metrangolo, G. Resnati. *Tetrahedron* **56** 5535–5550 (2000).
- [27] W. Li, L. Spada, N. Tasinato, S. Rampino, L. Evangelisti, A. Gualandì, P. G. Cozzi, S. Melandri, V. Barone, C. Puzzarini. *Angew. Chem. Int. Ed.* **57** 13853–13857 (2018).
- [28] P. Politzer, J. S. Murray. *Crystals* **9** 165 (2019).
- [29] I. Chircă, C. Silvestru, H. J. Breunig, C. I. Raț. *Inorg. Chim. Acta* **475** 155–160 (2018). CCDC no.: 1565715.
- [30] L. Dostál, I. Císařová, R. Jambor, A. Růžička, R. Jirásko, J. Holeček. *Organometallics*. **25**, 4366–4373 (2006). CCDC no.: 609167.
- [31] J. S. Murray, P. Lane, T. Clark, K. E. Riley, P. Politzer. *J. Mol. Model.* **18** 541–548 (2012).
- [32] V. Gutmann. *Coord. Chem. Rev.* **18**, 225–255 (1976).
- [33] L. O. Meller, D. Himmel, J. Stauffer, G. Steinfeld, J. Slattery, G. Santiso-Quiçones, V. Brecht, I. Crossing. *Angew. Chem. Int. Ed.* **47**, 7659–7663 (2008).
- [34] S. Scheiner. *Acc. Chem. Res.* **46**, 280–288 (2013).
- [35] J. S. Murray, P. Lane, P. Politzer. *Intern. J. Quantum Chem.* **107**, 2286–2292 (2007).
- [36] N. W. Alcock. *Adv. Inorg. Chem. Radiochem.* **15**, 1–58 (1972).
- [37] S. Scheiner. *J. Lu. Chem. Eur. J.* **24**, 8167–8177 (2018).
- [38] S. Scheiner. *Inorg. Chem.* **59**, 9315–9324 (2020).
- [39] Y. Mourad, A. Atmani, Y. Mugnier, H. J. Breunig, K. H. Ebert. *J. Organomet. Chem.* **476**, 47–54 (1994). CCDC no.: 1236532.
- [40] O. Mundt, G. Becker, H. Stadelmann, H. Thurn. *Z. Anorg. Allg. Chem.* **617**, 59–71 (1992). CCDC nos.: 1201650 and 1201651.
- [41] P. L. Millington, D. B. Sowerby. *J. Organomet. Chem.* **480**, 227–234 (1994). CCDC no.: 1237239.
- [42] A. Bauzá, A. V. Sharko, G. A. Senchyk, E. B. Rusanov, A. Frontera, K. V. Domasevitch. *CrystEngComm* **19** 1933–1937 (2017).
- [43] D. Walther, O. Klobes, M. Stollenz, W. Imhof, H. Gorts. *Cambridge Structural Database Communication, Private Communication*, 2003. CCDC no.: 114830.
- [44] D. Lu, M. L. Coote, J. Ho, N. L. Kilah, C.-Y. Lin, G. Salem, M. L. Weir, A. C. Willis, S. B. Wild, P. J. Dilda. *Z. Anorg. Allg. Chem.* **617**, 59–71 (1992). *Organometallics* **31**, 1808–1816 (2012). CCDC no.: 847338.
- [45] A.-J. Di Maio, A. L. Rheingold. *Organometallics* **10**, 3764–3766 (1991). CCDC no.: 1286040.
- [46] D. Rosmann, K.-W. Klinkhammer, A. Schmidt. *Monatsh. Chem.* **127**, 461–468 (1996). CCDC no.: 1248036.
- [47] N. Camerman, J. Trotter. *Can. J. Chem.* **41**, 460–464 (1963). CCDC no.: 1127915.
- [48] S. Solyntjes, J. Bader, B. Neumann, H.-G. Stammer, N. Ignatev, B. Hoge. *Chem. Eur. J.* **23**, 1557–1567 (2017). CCDC no.: 1474780.
- [49] J. W. Wielandt, S. Petrie, N. L. Kilah, A. C. Willis, R. D. Dewhurst, F. Belaj, A. Orthaber, R. Stranger, S. B. Wild. *Aust. J. Chem.* **69**, 524–533 (2016). CCDC no.: 1433855.
- [50] J. J. Weigand, K.-O. Feldmann, F. D. Henne. *J. Am. Chem. Soc.* **132** 16321–167323 (2010). CCDC no.: 782415.
- [51] F. D. Henne, A. T. Dickschat, F. Hennesdorf, K.-O. Feldmann, J. J. Weigand. *Inorg. Chem.* **54**, 6849–6861 (2015). CCDC no.: 1055388.
- [52] J. B. Waters, Q. Chen, T. A. Everitt, J. M. Goicoechea. *Dalton Trans.* **46**, 12053–12066 (2017). CCDC no.: 1560212.
- [53] E. Atrián-Blasco, S. Gascón, M. J. Rodríguez-Yoldi, M. Laguna, E. Cerrada. *Eur. J. Inorg. Chem.* 2791–2803 (2016). CCDC no.: 1454024.
- [54] Y. Uchiyama, R. Murakamia, J. Sugimoto. *Phosph. Sulfur Sil. Relat. Elem.* **190**, 633–637 (2015). CCDC nos.: 1019728 and 1019729.
- [55] S. S. Batsanov. *Inorg. Mat.* **37** 871–885 (2001).
- [56] P. F. Lang, B. C. Smith. *Dalton Trans.* **43** 8016–8025 (2014).
- [57] J. Lu, S. Scheiner. *Molecules*. **24** 2822 (2019).
- [58] S. Benz, A. I. Poblador-Bahamonde, N. Low-Ders, S. Matile. *Angew. Chem. Int. Ed.*, **57** 5408–5412 (2018).
- [59] M. S. Taylor. *Coord. Chem. Rev.* **413** 213270 (2020).
- [60] D. Scheiner. *Chem. Eur. J.* **22** 18850–18858 (2016).
- [61] P. R. Joshi, N. Ramanathan, K. Sundararajan, K. Sankaran. *J. Phys. Chem. A* **119**, 3440–3451 (2015).
- [62] P. K. Sruthi, N. Ramanathan, Shubhra Sarkar, K. Sundararajan. *Phys. Chem. Chem. Phys.* **20** 22058–22075 (2018).
- [63] P. R. Joshi, N. Ramanathan, K. Sundararajan, K. Sankaran. *J. Mol. Spectr.* **331** 44–52 (2017).
- [64] M. Yang, D. Tofan, C.-H. Chen, K. M. Jack, F. P. Gabbai. *Angew. Chem. Int. Ed.* **57**, 13868–13872 (2018),
- [65] C. Leroy, R. Johannson, D. L. Bryce. *J. Phys. Chem. A* **123**, 1030–1043 (2019).
- [66] P. Scilabra, G. Terraneo, A. Daolio, A. Baggioli, A. Famulari, C. Leroy, D. L. Bryce, G. Resnati. *Cryst. Growth Des.* **20**, 916–922 (2020).
- [67] R. Mokrai, J. Barrett, D. C. Apperley, A. S. Batsanov, Z. Benko, D. Heift. *Chem. Eur. J.* **25** 4017–4024 (2019).
- [68] G. G. Briand, N. Burford. *Adv. Inorg. Chem.* **50** 285–357 (2000).
- [69] J. P. M. Lommerse, A. J. Stone, R. Taylor, F. H. Allen. *J. Am. Chem. Soc.* **118** 3108–3116 (1996).
- [70] C. Trujillo, G. Sánchez-Sanz, I. Alkorta, J. Elguero. *New J. Chem.* **39** 6791–6802 (2015).
- [71] M. Krasowska, A.-M. Fritzsche, M. Mehring, A. A. Auer. *ChemPhysChem* **20**, 2539–2552 (2019).
- [72] V. M. Cangelosi, M. A. Pitt, W. J. Vickaryous, C. A. Allen. L. N. Zakharov, D. W. Johnson. *Cryst. Growth Des.* **10** 3531–3536 (2010).

- 1  
2  
3 [73] I.-S. Ke, F. P. Gabbai. *Inorg. Chem.* **52** 7145–7151 (2013).  
4 [74] J.-M. Lehn, M. Mascal, A. DeCian, J. Fischer. *J. Chem. Soc., Chem. Commun.* 479–481 (1990).  
5 [75] A. D. Legon. *Chem. Soc. Rev.* **22** 153–163 (1993).  
6 [76] K. Lisac, F. Topić, M. Arhangelskis, S. Cepić, P. A. Julien, C. W. Nickels, A. J. Morris, T. Friščić, D. Cinčić. *Nature Comm.*  
7 **10**, 61 (2019). CCDC nos.: 1850441, 1850434, 1856642.  
8 [77] S. Zahn, R. Frank, E. Hey-Hawkins, B. Kirchner. *Chem. Eur. J.* **17** 6034–6038 (2011).  
9 [78] A. Bauza, D. Quinñero, P. M. Deya, A. Frontera. *Phys. Chem. Chem. Phys.* **14** 14061–14066 (2012).  
10 [79] A. Varadwaj, P. R. Varadwaj, H. M. Marques, K. Yamashita. *Inorganics*, **10** 149 (2022).  
11 [80] G. Cavallo, P. Metrangolo, T. Pilati, G. Resnati, G. Terraneo. *Cryst. Growth Des.* **14** 2697–2702 (2014).  
12 [81] D. C. Ghosh, R. Biswas. *Int. J. Mol. Sci.* **2003**, **4**, 379–407 (2003).  
13 [82] W. Lindemann, R. Wogerbauer, P. Berger. *Zeit. Anorg. Allg. Chem.* **437** 155–158 (1977). CCDC no.: 1275802.  
14 [83] F. B. Alhanash, N. A. Barnes, A. K. Brisdon, S. M. Godfrey, R. G. Pritchard. *Dalton Trans.* **41**, 10211–10218 (2012). CCDC  
15 no.: 881803.  
16 [84] K. H. Ebert, R. E. Schulz, H. J. Breunig, C. Silvestru, I. Haiduc. *J. Organomet. Chem.* **470**, 93–98 (1994). CCDC no.: 1292824.  
17 [85] K. P. Bhabak, G. Mugesh. *Chem. Eur. J.* **15** 9846–9854 (2009). CCDC no.: 724870.  
18 [86] A. Nordheider, E. Hupf, B. A. Chalmers, F. R. Knight, M. Buhl, S. Mebs, L. Checinska, E. Lork, P. S. Camacho, S. E.  
19 Ashbrook, K. S. A. Arachchige, D. B. Cordes, A. M. Z. Slawin, J. Beckmann, J. D. Woollins. *Inorg. Chem.* **54**, 2435–2446  
20 (2015). CCDC no.: 1485019.  
21 [87] A. Ruzicka, Z. Padelkova, P. Svec, V. Pejchal, L. Ceslova, J. Holecek. *J. Organomet. Chem.* **732**, 47–57 (2013). CCDC no.:  
22 918470.  
23 [88] M. Korenkova, R. Jambor, Z. Ruzickova, L. Dostal. *Inorg. Chem. Commun.* **69** 28–30 (2016). CCDC no.: 1442753.  
24 [89] N. Auner, R. Probst, F. Hahn, E. Herdtweck. *J. Organomet. Chem.* **459** 25–41 (1993). CCDC no.: 1300159.  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60